Spatial patterns of white spruce (*Picea glauca*) in Eastern North America

Michael W. Wade

Spring Quarter 2016-2017

Thesis submitted in completion of Honors Senior Capstone requirements for the DePaul University Honors Program

Thesis Director: Jalene M. LaMontagne, Department of Biological Sciences, DePaul University

Faculty Reader: Jessica M. Vogt, Department of Environmental Sciences and Studies, DePaul University

Abstract

Many species are responding to climate change by adjusting their spatial distribution and altering their growth patterns, especially near range limits. We studied historic growth of white spruce over two spatial scales. We studied populations in Michigan, Ontario, and New Brunswick to examine growth trends over distances of >1500 km, and then local populations separated by a distance of 5 km in Michigan. Trends in growth were compared to mean temperatures across sites. We demonstrate the use of regional climate, influenced at local scales by topographical features (e.g. presence of large water bodies), to study the effects of climate change.

Table of Contents

Introduction1	
Methods3	
Results6	
Discussion8	
References Cited11	
Figures & Tables14	

Introduction

Global temperature increase resulting from climate change affects the growth and distribution of a large number of species, including many perennial plants (McKenney et al., 2007). For instance, in the boreal forest of North America, tree populations have responded positively to temperature increase over short periods of time, in terms of individual growth (Stinziano & Way, 2014) and temperature increase has been shown to foster treeline movement both northward latitudinally and upwards in altitude (Wilmking et al., 2004). High-latitude forests store up to 49% of the total carbon stored by forest ecosystems globally (Dixon et al., 1994), and these populations are important in mitigating the effects of climate change.

A pervasive boreal forest species in North America is white spruce (*Picea glauca*). Populations of this conifer species are found throughout much of Canada from the east coast to the west, and into northern Canada and Alaska (Figure 1). At high latitudes and at high elevation, annual growth rates of individual white spruce trees have increased in response to higher air and soil temperatures (Danby & Hik, 2006). However, precipitation plays an important role in tree growth as well. An increase in air temperature was found to negatively impact the radial growth of white spruce populations in Alaska; likely due to physiological stress resulting from moisture (usually available as runoff from melted snow), being less available due to the temperature increase (Barber & Juday, 2000).

Because the distribution of white spruce spans a large trans-continental spatial range, accurately describing the species' growth response to climate change is dependent on understanding how populations across specific regions respond to temperature and other environmental variables. Previous studies relating white spruce tree growth to weather conditions, such as temperature, precipitation, and humidity, have focused on sites at high altitudes (Barber & Juday, 2000), and/or at high latitudes (Wilmking et al., 2004; Danby & Hik, 2006), near the species' northern range limit, in regions such as Yukon and Alaska. This project described populations near the southern range limit, where growing seasons are substantially longer than at higher latitudes (Rowe, 1972), and water availability is often provided by rainfall, rather than snowmelt (Holman et al., 2012; Notaro et al., 2015).

Our study sought to describe the effects of growing season temperature on annual growth of white spruce individuals near the southern limit of the species' range. Specifically, we focused on the radial growth of individuals per year. Two spatial scales were studied. First, large spatial extent, comparing populations that are separated from one another by distances of 600-1600 km. A second, local area within one site was also studied: i) lakeshore sites and ii) inland sites. In this region, a lake effect results in an average difference in temperatures of up to 2 °C between an area directly adjacent to Lake Superior and an area 5 km inland (Appendix; Hinkel & Nelson, 2012).

We hypothesized that annual growth rates would vary based on the location of individuals. Regions with greater correlation between growing season temperatures were expected to yield white spruce individuals with more similar annual growth rates. Additionally, warmer growing season temperatures were expected to result in decreased annual growth, as each population was located in temperate regions where higher annual temperatures often coincide with decreased precipitation (Trenberth & Jones, 2007; Cong & Brady, 2012; Betts et al., 2014), and cold temperatures are not extreme enough to restrict growth of white spruce individuals in these areas (Nienstaedt & Zasada; Rowe, 1972).

Methods

Study sites

At the large spatial extent, sites were located in Michigan, Ontario, and New Brunswick (Figure 1). The Michigan study site was located northwest of Powell, Michigan (46°52'08.3"N, 87°50'24.6"W), and included land maintained by the Huron Mountain Wildlife Foundation. The region is largely undeveloped and is dominated by old-growth forests and small freshwater ponds and lakes. Annually, average temperatures range from $6 - 11^{\circ}$ C and the average rainfall is 81 cm (Weather Underground). The presence of Lake Superior results in inland areas being relatively warmer than those along the lakeshore (Hinkel & Nelson, 2012). The Ontario site was located near North Bay (46°18'58.2"N, 79°28'07.1"W), inland from Lake Nipissing. Annually, average temperatures range from $0 - 9^{\circ}$ C and the average rainfall is 105 cm (Gov. of Canada). The New Brunswick site was located in Lorneville, NB, which is approximately 20 km southwest of Saint John, NB (45°10'55.8"N, 66°09'20.3"W). The region is inland from the northern shore of the Bay of Fundy. Annually, average temperatures range from $0 - 10^{\circ}$ C and the average rainfall is 143 cm (Gov. of Canada).

Core collection & processing

White spruce trees were cored from sites near North Bay, Ontario and Lorneville, New Brunswick in 2016. Ten living trees were selected at each of these sites, and cores were collected at breast height (1.3 m) and stored in opaque plastic tubes for transportation and storage. In the lab, each core was transferred onto a core mount, rotated such that the rings faced directly upward, and secured with carpenter's glue. Once each core was secured in its mount, it was sanded down with 100, 150, and 250 grit sandpaper until a flat, smooth surface was formed

(Speer, 2010).

In Michigan, twenty-one white spruce cores were collected from six sites (three adjacent to Lake Superior ("lakeshore"), three approximately 5 km inland ("inland")) (Figure 2) in the Huron Mountains of Michigan in summer of 2012. Tree selection was based on diameter, with only the largest three adult living individuals at each site selected for coring. Cores were collected at 30 cm above ground level. Preparation of each core for processing was performed once more by securing each core in a core mount and sanding it down with 100, 150, and 250 grit sandpaper (Speer, 2010).

Data collection

Each core was viewed and each tree's radial growth increments were measured using a Velmex stage and dissecting microscope. These increments are indicated by distinguishable lines on the cores. MeasureJ2X software was used to record the width of rings from bark to pith. The measurements were checked for accuracy using COFECHA. Where dating errors were shown to be greater than 2-3 years, as reported by COFECHA, re-measurements of cores and adjustments using the program's autoregression feature were completed until the measurements were indicated as reasonably accurate (Grissino-Mayer, 2001). The corrected individual series were then detrended using ARSTAN, allowing age effects and general trends spanning multiple years to be removed from analysis (Cook et al., 1986). Finally, residual chronology indices were tabulated using ARSTAN. The outputs from ARSTAN were in raw growth increments or the indexed residual increments, which were each used for analysis.

Historic temperature data for each of the sites were collected from weather stations nearest to each site. For the Michigan site, historic monthly temperature means were provided by a weather station in Big Bay, MI (46°50'05.8"N, 87°42'42.6"W;

http://www.intellicast.com/Local/History.aspx?location=USMI0082). The Ontario temperature data were provided by a weather station at the North Bay Airport (46°21'25.6"N, 79°25'43.0"W; http://climate.weather.gc.ca/historical_data/search_historic_data_e.html), and the New Brunswick temperature data were provided by a weather station in Saint John, NB (45°15'54.9"N, 66°03'11.9"W; https://www.wunderground.com/history/airport/CYSJ). Mean temperatures were calculated from April 1 – September 31 of each year from the earliest year available to the year of core collection. The growing season length was based on previous data collected by the United States Forestry Service (Nienstaedt & Zasada, 1990).

Data analysis

Age effects on annual growth of individuals, which can result in the magnification of growth increments of older trees, were minimized by generating standardized indices for radial growth; annual growth residuals were used for analyses rather than raw ring width values (Cook & Peters, 1997). For the large spatial extent, comparing the growth of individuals between the New Brunswick, Ontario, and combined Michigan sites, annual residual growth was compared with mean growing season temperature at each site. To determine the spatial correlation between mean annual growing season temperature across distance classes of 500 km bin sizes, a modified, non-centered spatial correlogram was generated using the correlog.nc function in the NCF package in R version 3.3.1 GUI 1.68 for Mac OSX (Bjornstad, 2016). Similarly, a spatial correlogram was generated to determine the correlation between mean annual growth across distance classes of 500 km bin sizes.

For the local scale including only the Michigan sites, individual growth was divided by

diameter at breast height, so that a standardized measurement for annual growth could be used for analysis. This method of standardization was preferred over using the indices from before, so that long-term effects of the same regional temperature could be observed, while still accounting for age effects on growth. Mean growth per ring width was compared between the lakeshore and inland populations to determine whether proximity to Lake Superior influenced annual growth. Finally, a linear mixed model (ANCOVA) was used to test the effects of proximity to Lake Superior on the relationship between mean annual residual growth and mean regional growing season temperature. All analyses were performed in R version 3.3.1 GUI 1.68 for Mac OSX.

Results

Large spatial extent (up to 1500 km)

At the Michigan site, the median age of individuals was 45 years, with the oldest individual dating back to 1916 and the youngest to 1993. The average core length at the Michigan site was 180.7 \pm 63.5 mm. At the Ontario site, individuals were much younger, with the median age being 13.5 years. The oldest individuals dated back to 2000, and the youngest to 2007. The average core length at the Ontario site was 76.1 \pm 12.9 mm. Finally, at the New Brunswick site, the median age of individuals to 2006. The average core length at the New Brunswick site was 96.1 \pm 43.0 mm. The average core lengths were significantly different across each site (one-way ANOVA, df = 34, F = 17.3, P < 0.001).

Growing season temperatures across each site were significantly different (one-way ANOVA, df = 94, F = 26.2, P < 0.001), with New Brunswick being the warmest site and Michigan being the coolest site, on average. The mean regional growing season temperature at

the Michigan site was the coolest of the three sites at 11.5 ± 1.0 °C between 1970 and 2012. Annual residual growth was shown to have a negative relationship with mean growing season temperature at this site, but the relationship was not significant (R² = 0.01, df = 29, F = 1.6, P = 0.2) (Figure 3a & 3b). The mean regional growing season temperature at the Ontario site was 12.6 ± 0.5 °C between 1970 and 2016. The relationship between annual residual growth and growing season temperature was also negative at the Ontario site, and was not significant (R² = 0.003, df = 15, F = 0.04, P = 0.8) (Figure 3c & 3d). Finally, the mean regional growing season temperature at the New Brunswick site was 13.3 ± 0.9 °C between 1970 and 2016. The relationship between annual residual growing season temperature at the New Brunswick site was 13.3 ± 0.9 °C between 1970 and 2016. The relationship between 1970 and 2016. The season temperature at the New Brunswick site was 13.3 ± 0.9 °C between 1970 and 2016. The season temperature at the New Brunswick site was 13.3 ± 0.9 °C between 1970 and 2016. The season temperature at this site was positive, but the relationship was not significant (R² = 0.01, df = 45, F = 1.7, P = 0.2), (Figure 3e & 3f).

The spatial correlograms indicated that mean growing season temperature was positively and significantly correlated within the same site, while negatively and significantly correlated between the Michigan and Ontario sites (Figure 4a; separated by just over 600 km). The correlations between mean growing season temperature of sites separated by distances greater than 1000 km were not significant. (Figure 4b). A similar trend was observed when correlating mean residual growth of individuals at each site, though the scale in correlations was much smaller than for temperature. Individuals located within the same site had the greatest correlation between annual residual growth, and the correlation was significant, but was low in magnitude (Figure 4b). The correlation between residual growth of individuals at the Michigan site and those at the Ontario site was negative and significant. Correlations between individuals separated by distances greater than 1000 km were not significant.

Local scale (~5 km)

From 1970 to 1997, inland individuals in Michigan had a greater mean growth than lakeshore individuals for all but one year (Figure 5). Between 1998 and 2012, however, there were several years of greater mean growth at the lakeshore site. Over the full 42-year span, individuals at the lakeshore site had a mean annual growth per core length of 0.010 ± 0.004 mm, while individuals at the inland site had a mean annual growth per core length of 0.012 ± 0.003 mm. The difference in mean annual growth per diameter was significant (paired t-test, df = 42, T = -3.9, P < 0.001). Mean residual growth had a negative, non-significant relationship with mean annual growing season temperature for the region at both the lakeshore (R² = 0.06, df = 29, F = 1.7, P = 0.2) and inland (R² = 0.07, df = 28, F = 2.3, P = 0.1) sites (Figure 6). While the yintercept for the inland site was significantly higher than that of the lakeshore site (ANCOVA, est. = 1.6, T = 4.1, P = 0.0001), there was no significant effect of site on response to mean regional temperature (ANCOVA, est. = -0.13, T = -0.232, P = 0.8).

Discussion

For each of the large-extent sites, growing season temperature was shown to be a relatively weak predictor for residual growth of white spruce individuals. The Michigan site featured the strongest trend in relating annual growth to growing season temperature, with warmer temperatures resulting in relatively decreased growth in the same year. This relationship argues against the positive growth response of white spruce to warming temperatures, as presented in some prior studies (Danby & Hik, 2006), but reinforces the conflicting growth responses observed in other studies involving white spruce populations at more northern latitudes (Barber & Juday, 2000; Wilmking et al., 2004). In each of these studies, individual growth was

shown to change based on the primary limiting factor (either temperature or precipitation) within the studied populations' regions.

Individuals within the same local-scale sites had the highest correlation in mean annual growth, and the correlation was negative between individuals from the Michigan and Ontario sites, which were separated by a distance of just over 600 kilometers. However, the correlation between individuals within the same site was just under 0.1, indicating that a high degree of variation exists even over a very small distance scale. This variation could result from the density of the forest surrounding each individual, as boreal individuals in less-dense areas often have greater annual growth (He & Duncan, 2000). Additionally, tree growth can be affected by the types of vegetation present in the undergrowth (Callaway & Walker, 1997; Nilsson & Wardle, 2005), which can vary over short distances.

Previous temperature studies at our sites near Lake Superior showed that inland areas are warmer during the summer than those directly alongside the lakeshore (Appendix; Hinkel & Nelson, 2012). Therefore, if temperature were a substantially limiting factor in this region, white spruce populations further inland would be expected to grow more, on average, than the lakeshore populations, and both should indicate a positive growth response to regional temperature. While inland individuals were shown to have grown more than lakeshore individuals between 1970 and 2012, both populations indicated a negative growth response to temperature. Consequently, it was determined that temperature must not be the primary limiting factor for white spruce individuals in this region.

One potential explanation for why temperature is not limiting is the size of the individuals selected. Mature white spruce are relatively large, and able to compete effectively for access to direct sunlight. This allows individuals to be less limited by regional temperature

fluctuations, so long as the range in temperature is not extreme (Wilmking et al., 2004). Furthermore, we chose the largest individuals at each site, which emerge through the forest canopy and are therefore less likely to be limited by temperature than individuals occupying the undergrowth, where temperatures are cooler due to restricted sunlight (Louis et al., 2005). A much more plausible limiting factor for these individuals would be access to water. Temperature and precipitation are often inversely related; a warmer growing season likely corresponds with decreased precipitation in the same year (Trenberth & Jones, 2007; Cong & Brady, 2012; Betts et al., 2014). Because water availability is likely a primary limitation on growth of white spruce along the species' southern range limit (Goldstein et al., 1985), a negative relationship would be expected between temperature and annual growth of individuals.

At higher latitudes, where temperatures range into extreme lows, temperature is the primary limiting factor, and warmer growing seasons will benefit the growth of individuals (Danby & Hik, 2006). However, in regions with milder annual temperatures, precipitation is the primary limiting factor for growth of white spruce. Increasing temperatures resulting from climate change will likely result in decreased individual growth of individuals near the southern limit of the species' range, due to the negative relationship observed between temperature and precipitation within the region (Trenberth & Jones, 2007; Cong & Brady, 2012; Betts et al., 2014). Because of the increased growth expected at high latitudes and decreased growth expected at lower latitudes, the overall effect of warming temperatures could likely be a shift northward in the overall distribution of white spruce.

References cited

- Barber, V. A. & Juday, G. P. (2000). "Reduced growth of Alaskan white spruce in the twentieth century from temperature-induced stress." *Nature*, 405: 668-673.
- Betts, A. K., Desjardins, R., Worth, D., & Beckage, B. (2014). "Climate coupling between temperature, humidity, precipitation, and cloud cover over the Canadian Prairies." *Journal of Geophysical Research: Atmospheres*, 119: 13,305-13,326.
- Bjornstad, O. N. (2016). "ncf: Spatial Nonparametric Covariance Functions." R package version 1.1-7. <u>https://CRAN.R-project.org/package=ncf</u>.
- Bonan, G. B. (2008). "Forests and climate change: forcings, feedbacks, and the climate benefits of forests." *Science*, 320 (5882): 1444-1449.
- Callaway, R. M. & Walker, L. R. (1997). "Competition and facilitation: a synthetic approach to interactions in plant communities." *Ecology*, 78 (7): 1958-1965.
- Cong, R & Brady, M. (2012). "The Interdependence between Rainfall and Temperature: Copula Analyses." *The Scientific World Journal*, 2012: 1-11.
- Cook, E. R. & Adams, R. K, & Fritts, H. C. (1986). "User manual for program ARSTAN." in Tree-Ring Chronologies of Western North America: California, eastern Oregon and northern Great Basin, 50-65.
- Cook, E. R. & Peters, K. (1997). "Calculating unbiased tree-ring indices for the study of climatic and environmental change." *The Holocene*, 7 (3): 361-370.
- Danby, R. K. & Hik, D. S. (2006). "Responses of white spruce (*Picea glauca*) to experimental warming at a subarctic alpine treeline." *Global Change Biology*, 13 (2): 437-451.
- Dixon, R. K., Brown, S., Houghton, R. A., Solomon, A. M., Trexler, M. C., & Wisniewski, J. (1994). "Carbon pools and flux of global forest ecosystems." *Science*, 263: 185-190.
- Field, C. B., Beherenfeld, M. J., Randerson, J. T., & Falkowski, P. (1998). "Primary production of the biosphere: integrating terrestrial and oceanic components." *Science*, 281: 237-240.
- Goldstein, G. H., Brubaker, L. B., & Hinckley, T. M. (1985). "Water relations of white spruce (*Picea glauca* (Moench) Voss) at tree line in north central Alaska." *Canadian Journal of Forest Research*, 15 (6): 1080-1087.
- Government of Canada: Environmental and natural resources. "Past weather and climate." Online. http://climate.weather.gc.ca/historical_data/search_historic_data_e.html.

- Grissino-Mayer, H.D. (2001). "Evaluating crossdating accuracy: A manual and tutorial for the computer program COFECHA." *Tree-Ring Research*, 57 (2): 205-221.
- He, F., & Duncan, R. P. (2000). "Density-dependent effects on tree survival in an old-growth Douglas fir forest." *Ecology*, 88 (4): 676-688
- Hinkel, K. M. & Nelson, F. E. (2012). "Spatial and temporal aspects of the lake effect on the southern coast of Lake Superior." *Theoretical Applied Climatology*, 2012 (109): 415-428
- Holman, K. D., Gronewold, A., Notaro, M., & Zarrin, A. (2012). "Improving historical precipitation estimates over the Lake Superior Basin." *Geophy. Research Letters*, 39 (3).
- Intellicast. "Historic average: Big Bay, Michigan." Online. http://www.intellicast.com/Local/History.aspx?location=USMI0082.
- Louis, A., Ounis, A., Ducruet, J. M., Evian, S., Laurila, T., Thum, T., Aurela, M., Wingsle, G., Alonso, L., Pedros, R., & Moya, I. (2005). "Remote sensing of sunlight-induced chlorophyll fluorescence and reflectance of Scots pine in the boreal forest during spring recovery." *Remote Sensing of Environment*, 96: 37-48.
- McKenney, D. W., Pedlar, J. H., Lawrence, K., Campbell, K., & Hutchinson, M. F. (2007). "Potential impacts of climate change on the distribution of North American trees." *Bioscience*, 57 (11): 939-948.
- Nienstaedt, H. & Zasada, J. C. (1990). "*Picea glauca* ((Moench) Voss)." United States Forestry Service. Online. https://www.na.fs.fed.us/spfo/pubs/silvics_manual/Volume_1/picea/glauca.htm.
- Nilsson, M. C. & Wardle, D. A. (2005). "Understory vegetation as a forest ecosystem driver: evidence from the Northern Swedish Boreal Forest." *Frontiers in Ecology & the Environment*, 3 (8): 421-428.
- Notaro, M., Bennington, V., & Vavrus, S. (2015). "Dynamically downscaled projections of lakeeffect snow in the Great Lakes Basin." *Journal of Climate*, 28 (4): 1661.
- Nowak, D. J. & Crane, D. E. (2001). "Carbon storage and sequestration by urban trees in the USA." *Environmental Pollution*, 116: 381-389.
- Rowe, J. S. (1972). "Forest regions of Canada: boreal forest region." Canadian Forestry Service; Department of the Environment. pp. 14-64
- Speer, J. H. (2009). "Field and Laboratory Methods." *In: Fundamentals of Tree-Ring Research*. University of Arizona Press.

- Stinziano, J. R. & Way, D. A. (2014). "Combined effects of rising [CO₂] and temperature on boreal forests." *Botany*, 92: 425-436.
- Tebaldi, C., Smith, R. L., Nychka, D., & Mearns, L. O. (2005). "Quantifying uncertainty in projections of regional climate change: a Bayesian approach to the analysis of multimodel ensembles." *Journal of Climate*, 18: 1524-1540.
- Trenberth, K.E., & Jones, P.D. (2007). "Observations: Surface and Atmospheric Climate Change." In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Weather Underground. "Weather history for Saint John, New Brunswick." Online. https://www.wunderground.com/history/airport/CYSJ.
- Wilmking, M., Juday, G. P., Barber, V. A., & Zald, H. S. (2004). "Recent climate warming forces contrasting growth responses of white spruce at treeline in Alaska through temperature thresholds." *Global Change Biology*, 10: 1724-1736.

Figures



Figure 1. Range of *Picea glauca* distribution, with locations of study sites indicated.



Figure 2. Michigan site map. Sites 1, 2, and 3 are inland, while sites 4, 5, and 6 are near the lakeshore.



Figure 3. Mean residual growth of white spruce and mean growing season temperatures at the (a) Michigan, (c) Ontario, & (e) New Brunswick sites. Residual growth of white spruce vs mean growing season temperatures for the (b) Michigan, (d) Ontario, & (f) New Brunswick sites.



Figure 4. (a) Correlation of mean growing season temperature over distance class. (b) Correlation of mean residual growth over distance class. Note change in scale on y-axis.



Figure 5. Difference in mean growth of white spruce per diameter between inland and lakeshore sites. Positive values indicate growth at inland sites was greater than at lakeshore sites. Negative values indicate growth at lakeshore site was greater than at inland sites.



Figure 6. Mean residual growth of white spruce at lakeshore and inland sites vs. regional growing season temperature from 1970-2012.

Appendix 1



Difference in monthly temperature between inland and shoreline near Lake Superior. Positive values indicate inland site was warmer than lakeshore site. From Hinkel & Nelson, 2012.